

MK2_A for Immunoprecipitation of MK2 from mammalian cell extract

Only for research applications, not for diagnostic or therapeutic use.

Introduction MK2 belongs to the family of serine/threonine kinases. In response to cellular stress it is phosphorylated and activated by MAP kinase p38. MK2-Trap is excellent for fast and efficient one-step isolation of MK2 and its interacting factors from cellular extract. Isolated MK2 protein may be used further for immunoblot analysis, mass spectrometry, and kinase assays. MK2-Trap utilizes small recombinant alpaca antibody fragments covalently coupled to the surface of agarose beads.

Specificity Species-Reactivity: tested on human, mouse, hamster
Posttranslational Modifications: MK2-Trap recognizes unphosphorylated MK2 and Phospho-MK2 (Thr222). Specificity on Phospho-MK2 (Thr334) was not tested.

Content	Reagent	Code	Quantity
	MK2-Trap_A	mta-20	20 reactions (0.5 ml slurry)
	MK2-Trap_A	mta -100	100 reactions (2.5 ml slurry)
	MK2-Trap_A	mta -200	200 reactions (5 ml slurry)
	MK2-Trap_A	mta -400	400 reactions (10 ml slurry)

Bead properties Bead size: ~ 90 µm (cross-linked 4% agarose beads)
Storage buffer: 20% EtOH

Stability and Storage Shipped at ambient temperature. Upon receipt store at +4°C.
Stable for 1 year. Do not freeze.

Required solutions **Suggested buffer composition**

Buffer	Composition
Lysis buffer (CoIP)	10 mM Tris/Cl pH 7.5; 150 mM NaCl; 0.5 mM EDTA; 0.5% NP-40
10x RIPA buffer	100 mM Tris/Cl pH 7.5; 1,5 M NaCl; 50 mM EDTA; 1% SDS; 10% Triton X-100; 10% Deoxycholate
Dilution/Wash buffer	10 mM Tris/Cl pH 7.5; 150 mM NaCl; 0.5 mM EDTA
Glycine-elution buffer	200 mM glycine pH 2.5
2 x SDS-sample buffer	120 mM Tris/Cl pH 6.8; 20% glycerol; 4% SDS, 0.04% bromophenol blue; 10% β-mercaptoethanol

Related products	MK2 Toolbox	Code
	MK2-Trap protein	mt-250
	MK2-Trap_A Kit	mtak-20
	Blocked agarose beads	bab-20
	Spin columns	sct-10; sct-20; sct-50

Support Please refer to our FAQ section at www.chromotek.com or contact support@chromotek.com

Protocol for Immunoprecipitation of MK2-Fusion Proteins using MK2-Trap_A

Harvest cells

For one immunoprecipitation reaction the use of $\sim 10^6 - 10^7$ mammalian cells (approx. one 10-cm dish) is recommended. To harvest adherent cells, aspirate growth medium, add 1 ml ice-cold PBS to cells and scrape cells from dish. Transfer cells to a pre-cooled tube, spin at 500 g for 3 min at +4°C and discard supernatant. Wash cell pellet twice with ice-cold PBS, gently resuspending the cells. After washing:

Lyse cells

1. Resuspend cell pellet in 200 μ l ice-cold lysis buffer by pipetting or using a syringe.
note: Supplement lysis buffer with protease inhibitors and 1 mM PMSF (not included).
optional for nuclear/chromatin proteins: Use RIPA buffer supplemented with 1 mg/ml DNase, 2.5 mM MgCl₂, protease inhibitors and 1 mM PMSF (not included).
2. Place the tube on ice for 30 min with extensively pipetting every 10 min.
3. Centrifuge cell lysate at 20.000x g for 10 min at +4°C. Transfer lysate to a pre-cooled tube. Add 300 μ l dilution buffer to lysate. Discard pellet.
note: At this point cell lysate may be put at -80°C for long-term storage.
optional: Add 1 mM PMSF and protease inhibitors (not included) to dilution buffer.
We recommend that during incubation with the beads the final concentration of detergents does not exceed 0.2% to avoid unspecific binding to the matrix. If required, use more dilution buffer to dilute the supernatant accordingly.

Equilibrate beads

4. Vortex MK2_A beads and pipette 25 μ l bead slurry into 500 μ l ice-cold dilution buffer. Centrifuge at 2.500x g for 2 min at +4°C. Discard supernatant and repeat wash twice.

Bind proteins

5. Add diluted lysate (step 3) to equilibrated MK2_A beads (step 4). If required, save 50 μ l of diluted lysate for immunoblot analysis. Tumble end-over-end for 1 hour at 4°C.
6. Centrifuge at 2.500x g for 2 min at +4°C. If required, save 50 μ l supernatant for immunoblot analysis. Discard remaining supernatant.

Wash beads

7. Resuspend MK2_A beads in 500 μ l ice-cold dilution buffer. Centrifuge at 2.500x g for 2 min at +4°C. Discard supernatant and repeat wash twice.
optional: Increase salt concentration in the second washing step up to 500 mM.

Elute proteins

8. Resuspend MK2_A beads in 100 μ l 2x SDS-sample buffer.
9. Boil resuspended MK2_A beads for 10 min at 95°C to dissociate immunocomplexes from MK2_A beads. MK2_A beads can be collected by centrifugation at 2.500x g for 2 min at 4°C and SDS-PAGE is performed with the supernatant.
10. *optional instead of steps 8 and 9: elute bound proteins by adding 50 μ l 0.2 M glycine pH 2.5 (incubation time: 30 sec under constant mixing) followed by centrifugation. Transfer the supernatant to a new tube and add 5 μ l 1M Tris base pH 10.4 for neutralization. To increase elution efficiency this step can be repeated.*

Limited Use Label License

The purchase of this product conveys to the buyer the limited, non-transferable right to use the purchased amount of the product and components of the product to perform internal research for the sole benefit of the purchaser. No right to resell this product or any of its components is conveyed expressly or by implication. For information on obtaining additional rights, please contact licensing@chromotek.com.