

# V5-Trap™ Agarose

Product code: v5ta



---

## Introduction

The ChromoTek V5-Trap™ Agarose consists of an anti-V5-tag Nanobody (VHH), which is covalently bound to agarose beads. V5-Trap Agarose is used to immunoprecipitate V5-tagged fusion proteins from cell extracts of various organisms like mammals, plants, bacteria, yeast, insects etc.

## Properties

**Ligand:** Anti-V5-tag single domain antibody fragment (VHH, Nanobody)

**Reactivity:** Specifically binds to V5-tag sequence (GKPIPPLLGLDST). Compatible with N-, C-terminal or internal tagging.

**Binding capacity:** 15 µg of recombinant V5-tagged protein (~30 kDa) per 25 µL bead slurry

**Bead size:** 90 µm (cross-linked 4 % agarose beads)

**Buffer compatibility:** See *Wash buffer compatibility table*.

**Storage buffer:** 20 % ethanol

**Storage conditions:** Upon receipt store at +4°C. Do not freeze!

**Stability:** Stable for 1 year upon receipt.

**Shipment:** Shipped at ambient temperature.

**RRID:** AB\_2868497

## Suggested buffer compositions

### Required buffer solutions

*NEW: Update of Wash buffer components.*

Buffer	Composition
Lysis buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5 % Nonidet™ P40 Substitute (adjust the pH at +4°C)
RIPA buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.1 % SDS, 1 % Triton™ X-100, 1 % deoxycholate (adjust the pH at +4°C)
Dilution buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA (adjust the pH at +4°C)
Wash buffer	10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.05 % Nonidet™ P40 Substitute, 0.5 mM EDTA (adjust the pH at +4°C)
2x SDS-sample buffer	120 mM Tris/Cl pH 6.8, 20 % glycerol, 4 % SDS, 0.04 % bromophenol blue, 10 % β-mercaptoethanol
Acidic elution buffer	100 mM glycine pH 2.0 (adjust the pH at +4°C)
Neutralization buffer	1 M Tris pH 10.4 (adjust the pH at +4°C)
Peptide elution buffer	2 mg/mL V5-peptide (v5p-1) reconstituted in PBS

*Note: Use your equivalent cell lysis buffer for other cell types like yeast, plants, insects, bacteria.*

*Note: Consider using a Wash buffer without detergent for co-immunoprecipitation.*

*Note: Use Peptide elution buffer or Acidic elution buffer for elution under native conditions.*

# V5-Trap™ Agarose

Product code: v5ta



## Wash buffer compatibility table

Buffer ingredients	Max. concentration
β-mercaptoethanol	10 mM
DTT	5 mM
NaCl	1 M
Nonidet™ P40 Substitute	tested up to 2 %
SDS	0 %
TCEP	5 mM
Triton™ X-100	tested up to 2 %
Urea	4 M

## Product sizes

Product	Product code	Size
V5-Trap™ Agarose	v5ta-10	10 reactions (250 µL slurry)
	v5ta-20	20 reactions (500 µL slurry)
	v5ta-100	100 reactions (2.5 mL slurry)
	v5ta-200	200 reactions (5 mL slurry)
	v5ta-400	400 reactions (10 mL slurry)
V5-Trap™ Agarose Kit	v5tak-20	20 reactions (500 µL slurry) including buffers

## Immunoprecipitation protocol

### Cell material

The following protocol describes the preparation of mammalian cell lysate!

For other type of cells, we recommend using 500 µg of cell extract and start the protocol with step *Bead equilibration*.

### Mammalian cell lysis

Note: Harvesting of cells and cell lysis should be performed with ice-cold buffers. We strongly recommend to add protease inhibitors to the Lysis buffer to prevent degradation of your target protein and its binding partners.

For one immunoprecipitation reaction, we recommend using  $\sim 10^6$ - $10^7$  cells.

#### 1. Choice of lysis buffer:

- For cytoplasmic proteins, resuspend the cell pellet in 200 µL ice-cold Lysis buffer by pipetting up and down. Supplement Lysis buffer with protease inhibitor cocktail and 1 mM PMSF (not included).
- For nuclear/chromatin proteins, resuspend cell pellet in 200 µL ice-cold RIPA buffer supplemented with DNaseI (f.c. 75-150 Kunitz U/mL), MgCl<sub>2</sub> (f.c. 2.5 mM), protease inhibitor cocktail and PMSF (f.c. 1 mM) (not included).

2. Place the tube on ice for 30 min and extensively pipette the suspension every 10 min.

3. Centrifuge cell lysate at 17,000x g for 10 min at +4°C. Transfer cleared lysate (supernatant) to a pre-cooled tube and add 300 µL Dilution buffer supplemented with 1 mM PMSF and protease inhibitor cocktail (not included). If required, save 50 µL of diluted lysate for further analysis (input fraction).

### Bead equilibration

1. Resuspend the beads by gently pipetting up and down or by inverting the tube. Do not vortex the beads!
2. Transfer 25 µL of bead slurry into a 1.5 mL reaction tube.
3. Add 500 µL ice-cold Dilution buffer.
4. Sediment the beads by centrifugation at 2,500x g for 5 min at +4°C. Discard the supernatant.

*Note:* Alternatively, Spin columns (sct-10; -20; -50) can be used to equilibrate the beads.

### Protein binding

1. Add diluted lysate to the equilibrated beads.
2. Rotate end-over-end for 1 hour at +4°C.

## Washing

1. Sediment the beads by centrifugation at 2,500x g for 5 min at +4°C.
2. If required, save 50 µL of supernatant for further analysis (flow-through/non-bound fraction).
3. Discard remaining supernatant.
4. Resuspend beads in 500 µL Wash buffer.
5. Sediment the beads by centrifugation at 2,500x g for 5 min at +4°C. Discard remaining supernatant.
6. Repeat this step at least twice.
7. During the last washing step, transfer the beads to a new tube.

*Optional:* To increase stringency of the Wash buffer, test various salt concentrations e.g. 150-500 mM, and/or add a non-ionic detergent e.g. Triton™ X-100 (see *Wash buffer compatibility table* for maximal concentrations).

*Note:* Alternatively, Spin columns (sct-10; -20; -50) can be used to wash the beads.

## Elution with 2x SDS-sample buffer (Laemmli)

1. Remove the remaining supernatant.
2. Resuspend beads in 80 µL 2x SDS-sample buffer.
3. Boil beads for 5 min at +95°C to dissociate immunocomplexes from beads.
4. Sediment the beads by centrifugation at 2,500x g for 2 min at +4°C.
5. Analyze the supernatant in SDS-PAGE / Western Blot.

## Elution with Acidic elution buffer

1. Remove the remaining supernatant.
2. Add 50-100 µL Acidic elution buffer and constantly pipette up and down for 30-60 sec at +4°C or room temperature.
3. Sediment the beads by centrifugation at 2,500x g for 2 min at +4°C.
4. Transfer the supernatant to a new tube.
5. Immediately neutralize the eluate fraction with 5-10 µL Neutralization buffer.
6. Repeat this step at least once to increase elution efficiency.

*Note:* Alternatively, Spin columns (sct-10; -20; -50) can be used to separate the beads.

## Elution with Peptide elution buffer

1. Reconstitute V5-peptide in PBS to a final concentration of 2 mg/mL (1.4 mM). Vortex for a few minutes to dissolve the powder.
2. Dilute V5-peptide stock to 0.72 mg/mL (0.5 mM) in PBS.
3. Remove the remaining supernatant from the beads.
4. Add 100 µL of the diluted V5-peptide and mix with a pipette.
5. Mix for 10-20 min at room temperature or for 30-60 min at +4°C.

# V5-Trap™ Agarose

Product code: v5ta

---



6. Sediment the beads by centrifugation at 2,500x g for 2 min at +4°C.
7. Transfer the supernatant to a new tube.
8. Repeat this step at least once to increase elution efficiency.

*Note:* Alternatively, Spin columns (sct-10; -20; -50) can be used to separate the beads.

# V5-Trap™ Agarose

Product code: v5ta



## Product overview and related products

V5-tag toolbox	Product code
V5-Trap™ Agarose	v5ta-10; -20; -100
V5-Trap™ Agarose Kit	v5tak-20
V5-Trap™ Magnetic Agarose	v5tma-10; -20; -100
V5-Trap™ Magnetic Agarose Kit	v5tmak-20
iST V5-Trap™ Kit for IP/MS	v5tak-IST-8
Binding Control Agarose	bab-20
Binding Control Magnetic Agarose	bmab-20
Spin columns	sct-10; sct-20; sct-50
V5-peptide	v5p-1
V5 VHH, recombinant binding protein	v5t-250
V5-tag antibody [SV5-P-K], monoclonal mouse IgG1 kappa	v5ab-20; -200
Nano-Secondary® alpaca anti-mouse IgG1, recombinant VHH, Alexa Fluor® 488 [CTK0103, CTK0104]	sms1AF488-1-10; -100
Nano-Secondary® alpaca anti-mouse IgG1, recombinant VHH, Alexa Fluor® 568 [CTK0103, CTK0104]	sms1AF568-1-10; -100
Nano-Secondary® alpaca anti-mouse IgG1, recombinant VHH, Alexa Fluor® 647 [CTK0103, CTK0104]	sms1AF657-1-10; -100

For product details, information, and ordering visit [www.chromotek.com](http://www.chromotek.com).

# V5-Trap™ Agarose

Product code: v5ta



---

## Contact

support@chromotek.com

ChromoTek GmbH  
Am Klopferspitz 19  
82152 Planegg-Martinsried  
Germany  
phone: +49 89 124 148 80  
fax: +49 89 124 148 811

ChromoTek Inc.  
62-64 Enter Lane  
Islandia, NY 11749  
USA  
phone: 631 501 1058  
fax: 631 501 1060

## Disclaimer

Only for research applications, not for diagnostic or therapeutic use!

ChromoTek and GFP-Trap, RFP-Trap, Myc-Trap, Spot-Trap, Spot-Tag, Spot-Label, Spot-Cap, Nano-Secondary, F2H Kit, and Chromobody are registered trademarks of ChromoTek GmbH and Nano-CaptureLigand and V5-Trap are trademarks of ChromoTek GmbH. Nanobody is a registered trademark of Ablynx, a Sanofi company. Alexa Fluor is a registered trademark of Life Technologies Corporation, a part of Thermo Fisher Scientific Inc. Dynabeads is a trademark of Life Technologies AS, a part of Thermo Fisher Scientific Inc. SNAP-tag is a registered trademark and CLIP-tag is a trademark of New England Biolabs, Inc. Octet is a registered trademark of ForéBio, a Sartorius brand. Other suppliers' products may be trademarks or registered trademarks of the corresponding supplier each. Statements on other suppliers' products are given according to our best knowledge.